

Uplift History of Syenite Rocks of the Sushina Hill, Tamar Porapahar Shear Zone (TPSZ), Purulia: Constraints from Fission-track Ages of Two Cogenetic Minerals

Amal Kumar Ghosh*, Virendra Kumar Sharma and Rajeev Kumar Singh

Bhagwant University, Ajmer, Rajasthan-305004, India

Abstract

Fission-track ages of cogenetic minerals namely apatite, and zircon from syenite rocks of the Sushina hill, Tamar Porapahar Shear Zone (TPSZ), coupled with the corresponding closure temperatures of the minerals have been used to reveal the uplift history of syenite rocks. Offset of their FT ages indicates that the samples uplifted at the rate of 9.97 m/Ma during the period 535 Ma-970 Ma.

Keywords: Metamorphic minerals; Cogenetic minerals; Syenite rocks; Uranium samples

Introduction

It is widely accepted that radiometric ages determined on metamorphic minerals from orogenic belts reflect their cooling history rather than their primary crystallization [1-5]. Cooling histories obtained using different radiometric techniques on cogenetic minerals from a single sample often include zircon and apatite fission track ages as low-temperature bounds to a temperature interval of several hundred degrees [6-15].

Fission-track ages were determined on cogenetic minerals from syenite rocks of the Sushina hill, TPSZ. Sushina hill in TPSZ lies within Singhbhum Group (SG) of rocks. The North Singhbhum Mobile Belt (NSMB) in its northern margin has a tectonic boundary with the Chhotanagpur Gneissic Complex (CGC) along the TPSZ [16-22]. Singhbhum Shear Zone (SSZ) is located 40 km south of the TPSZ. At places, TPSZ passes either through rocks of SG or CGC. Singhbhum orogenic cycle experienced three major phases of deformation. CGC underwent four phases of deformation [23-28]. TPSZ witnessed reactivation near 500 Ma due to overthrusting and suffered rapid exhumation near 600 Ma [29-34]. TPSZ was thus affected by the intense deformation in the area adjoining this shear zone. With this background of geological events, it was thought that fission track dating on cogenetic minerals might help unravel the uplift history of syenite rocks of the Sushina hill.

Experimental Procedure

The samples for this study were processed in the laboratory of the Geological Survey of India, Kolkata, after obtaining permission from the Director General, GSI, Kolkata, West Bengal. The samples were prepared using standard separation, grinding and polishing techniques [35]. All the samples were prepared for the external detector method. AFT mounts were etched with 70% HNO₃ at room temperature for 30 s. Zircons were mounted in PFA Teflon. Zircons were etched in KOH-NaOH eutectic etchant [36-39] at 215°C on Spinot digital hot plate for ~8 hrs. The sample was placed in 48% HF for 2 hrs to clean up grains. After etching, mica sheets were firmly attached on the sample mounts. The samples were irradiated in the thermal facilities of FRMII at Garching, Germany together with dosimeter glass IRMM-540R (15 ppm). Mica sheets were etched using 48% HF at room temperature for 19 min [40-43]. The fission tracks were counted under a total magnification of 1000X. The calibrated area of one grid is 0.64×10⁻⁶ cm². Durango apatites were used as the age standard mineral, which was provided by Prof. Barry Paul Kohn, University of Melbourne,

Australia. FT age of zircon was determined using equation without zeta value. Zeta calibration was not performed on zircon because of the unavailability of age standard minerals [44-48].

Interpretation and Results

Fission track age determinations were made on 15apatite, and 18 zircon separates from syenite rocks from the Sushina hill shown in Figure 1. The mean fission track ages of apatite and zircon are 535.25 Ma, and 970 Ma respectively as shown in Table 1.

The uplift rate has been calculated according to the equation:

$$\text{Uplift rate} = \frac{\text{Cooling rate}}{\text{Geothermal gradient}} \quad (1)$$

Where, Cooling rate = $(T_1 - T_2) \div (A_1 - A_2)$

T_1, T_2 = Closure temperatures of cogenetic minerals

A_1, A_2 = Mean FT ages of cogenetic minerals

Average geothermal gradient of the order of 30°C/km has been adopted. Closure temperatures for apatite and zircon have been adopted 110°C and 240°C respectively.

FT age of apatite sample namely SAP has been calculated according to the equation [49]:

$$T = 1/\lambda_d \ln \{ 1 + \lambda_d Z G \rho_d \rho_s / \rho_i \} \quad (2)$$

Where, ρ_s is the surface density of etched spontaneous fission tracks, ρ_i is the surface density of etched induced fission tracks, and G is the integrated geometry factor of etched surface. $\lambda_d = 1.55 \times 10^{-10} \text{ yr}^{-1}$ = total decay constant of ²³⁸U, Z = calibration factor based on EDM of fission-track age standards. ρ_d = induced fission-track density for a uranium standard corresponding to the sample position during neutron irradiation.

*Corresponding author: Amal Kumar Ghosh, Ph.D. Student, Bhagwant University, Ajmer, Rajasthan-305004, India, Tel: + 9830520035; E-mail: ghosh.971@gmail.com

Received March 08, 2016; Accepted April 25, 2016; Published April 29, 2016

Citation: Ghosh AK, Sharma VK, Singh RK (2016) Uplift History of Syenite Rocks of the Sushina Hill, Tamar Porapahar Shear Zone (TPSZ), Purulia: Constraints from Fission-track Ages of Two Cogenetic Minerals. J Geol Geophys 5: 245. doi:10.4172/2381-8719.1000245

Copyright: © 2016 Ghosh AK, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Location map of the Sushina hill.

Sample name	Rock type and Mineral	$P_d \times 10^6$	N_d	$P_s \times 10^6$	N_s	$P_i \times 10^6$	N_i	$P\{x\}2(\%)$	U(ppm)	Mean age ± 16 (Ma)	No. of grain
SAP	Syenite, Apatite	1.925	1232	0.325	156	0.146	70	0.33	0.68	535.25 \pm 14.52	15
SZR	Syenite, Zircon	1.82	1168	17.66	452	1.773	244	15.39	41.65	970 \pm 7.94	18

Table 1: Results of AFT analyses : ages calculated using dosimeter glass IRMM-540R with 15 ppm U, zeta = 250, irradiated at FRMII, calibrated by traditional zeta approach and external detector method for apatite sample SAP, N=Number of grains, ρ – track densities given in 10^6 tr cm^{-2} , ρ_d – dosimeter track density, N_d – number of tracks counted on dosimeter, $\rho_s(\rho_i)$ – spontaneous (induced) track densities, $N_s(N_i)$ – number of counted spontaneous (induced) tracks, $P\{x\}$ – probability for obtaining 2x value for n degrees of freedom, where n=no. of grain – 1, Neutron flux(Φ) for zircon sample SZR = 1.75×10^{15} neutron/ cm^2 .

Region	Closure temperatures	Time Span (Ma)	Cooling rate ($^{\circ}C/Ma$)	Uplift rate (m/Ma)
Sushina hill in TPSZ	110 $^{\circ}C$ for apatite 240 $^{\circ}C$ for zircon	535-970	0.299	9.97

Table 2: Cooling and uplift rate of the Sushina hill.

FT age of zircon sample namely SZR has been calculated according to the equation without zeta value [50]:

$$T = 1 / \lambda_d \ln \left(1 + 9.25 \times 10^{-18} \rho_s / \rho_i \Phi \right) \text{ years} \quad (3)$$

Where, Φ = Neutron flux

The large age errors, e.g. 14.52% are found in sample SAP. As already known, low Uranium samples present a problem because of low induced track densities [51-55]. The P (X^2) test was performed to measure the uranium variation in the samples. A value of P(X^2) larger than 5% means that the grains are assumed to be a single age. Sample SAP failed the X^2 test, which may indicate bimodal distributions for the sample.

By applying apatite fission track analysis, the possible uplift rate of the Sushina hill was attempted to be revealed, which could be reflected by an offset age of two cogenetic minerals (Table 2).

Conclusion

The largest age error (14.52%) occurs in sample SAP. This high error is most likely due to a very low uranium concentration (0.68 ppm). As already known, low uranium samples place limits on how robust the ages could be. In low uranium samples, an exact match between the areas counted in the grains and the mica is often hard to achieve. An adjustment by eye is difficult and subjective because the outline of the induced tracks on the mica does not reflect the shape of the analyzed grain. In this study, closure temperatures for apatite and zircon have been adopted 110 $^{\circ}C$ and 240 $^{\circ}C$ respectively. In reality, the closure temperature concept cannot be straight forward applied.

For determining zircon FT age, zeta calibration was not performed. It places limit on precise calculation of FT age. Syenite rocks of the Sushina hill in TPSZ uplifted at the rate of 9.97 m/Ma in the range from 535 Ma-970 Ma.

Acknowledgement

I thank Prof. Barry Paul Kohn, "University of Melbourne", Australia for his overall support and advice throughout my time as a Ph.D. student. He also sent me age standard minerals (Durango apatite) for zeta calibration, free of charge. Without his contribution, my work would have never been possible.

I thank Prof. Richard Ketcham "University of Texas", U.S.A., who kindly reviewed my AFT models and provided valuable guidance for the improvement of the models.

I am highly indebted to the Director General of Geological Survey of India, 27, J.L. Nehru Road, Kolkata – 700 016, for his kind permission to perform my work in the laboratory of G.S.I, Kolkata.

I thank Mr. Partha Nag, Officer-in-Charge, WBMTDC, Purulia for his dedicated help with the field work.

I thank Mr. T. Ray Barman, Ex-Scientist, G.S.I, Kolkata, for his constructive advice.

Many thanks are due to the entire family of G.S.I., Kolkata, for their help and encouragement.

I thank the entire team of FRMII, Garching, Germany for providing me with use of the irradiation facility, free of charge.

References

- Acharya A, Basu SK, Bhaduri SK, Chaudhury BK, Ray S, et al. (2006) Proterozoic Rock Suites along South Purulia Shear Zone, Eastern India; Evidence for Rift-Related Setting. *Geological Society of India* 68: 1069-1086.
- Basu SK (1993) Alkaline-Carbonatite Complex in Precambrian of South Purulia Shear Zone, Eastern India; Its Characteristics and Minerals Potentialities. *Indian Minerals* 47: 179-194.
- Belton DX, Brown RW, Gleadow AJW, Kohn BP (2002) Fission Track Dating of Phosphate minerals and the Thermochronology of Apatite 16: 579-630.
- Brown R, Gallagher K, Johnson C (1998) Fission Track Analysis and its Applications to Geological Problems. *Annual Review Earth Planet Science* 26: 519-572.
- Carter A, Cliff PD, Dorr N, Gee DG, Liskar F, et al. (2012) Late Mesozoic – Cenozoic exhumation history of northern Svalbard and its regional significance; constraints from apatite fission track analysis. *Tectonophysics* 514-517.
- Dodge FCW, Ross DC (1971) Coexisting hornblendes and biotites from granite rocks near the San Andreas fault, California. *Journal of Geology* 79: 158-172.
- Fayon A, Peacock SM, Stump E, Reynolds SJ (2000) Fission track analysis of the footwall of the Catalina detachment fault, Arizona: Tectonic denudation, magmatism, and erosion. *J Geophys Res* 105: 11047-11062.
- Feinstein S, Kohn BP, Eyal M (1989) Significance of combined vitrinite reflectance and fission-track studies in evaluating thermal history of sedimentary basins: an example from southern Israel. In: Naeser ND, McCulloch TH (eds.) *Thermal History of Sedimentary Basins: Methods and Case Histories*. Springer-Verlag, Berlin pp: 197-216.
- Fitzgerald PG (1994) Thermochronological constraints on the post-Paleozoic tectonic evolution of the central Transantarctic Mountains, Antarctica. *Tectonics* 13: 818-836.
- Fitzgerald P, Gleadow AJW (1990) New approaches in fission track geochronology as a tectonic tool: Examples from the Transantarctic Mountains. *Nucl Tracks Radiat Meas* 17: 351-357.
- Fitzgerald PG, Sandiford M, Barrett PJ, Gleadow AJW (1986) Asymmetric extension in the Transantarctic Mountains and Ross Embayment. *Earth Planet Sci Lett* 86: 67-78.
- Fitzgerald PG, Fryxel JE, Wernicke BP (1991) Miocene crustal extension and uplift in southeastern Nevada: Constraints from fission track analysis. *Geology* 19: 1013-1016.
- Fitzgerald PG, Reynolds SJ, Stump E, Foster DA, Gleadow AJW (1993) Thermochronologic evidence for timing of denudation and rate of crustal extension of the South Mountain metamorphic core complex and Sierra Estrella, Arizona. *Nucl Tracks* 21: 555-563.
- Fitzgerald PG, Sorkhabi RB, Redfield TF, Stump E (1995) Uplift and denudation of the central Alaska Range: a case study in the use of apatite fission track thermochronology to determine absolute uplift parameters. *J Geophys Res* 100: 20175-20191.
- Fitzgerald PG, Munoz JA, Coney PJ, Baldwin SL (1999) Asymmetric exhumation across the Pyrenean orogen: implications for the tectonic evolution of a collisional orogen. *Earth Planet Sci Lett* 173: 157-170.
- Fleischer RL, Hart HR (1972) Fission track dating: Techniques and problems. In: Bishop WW, Miller DA, Cole S (eds.) *Calibration of hominid evolution*. Scottish Academic Press, Edinburgh pp: 135-170.
- Fleischer RL, Price PB (1964) Techniques for geological dating of minerals by chemical etching of fission fragment tracks. *Geochim Cosmochim Acta* 28: 1705-1715.
- Fleischer RL, Price PB, Symes EM (1964) On the origin of anomalous etch figures in minerals. *Am Mineral* 49: 794-800.
- Fleischer RL, Price PB, Walker RM (1965a) Effects of temperature, pressure and ionization of the formation and stability of fission tracks in minerals and glasses. *J Geophys Res* 70: 1497-1502.
- Fleischer RL, Price PB, Walker RM (1965b) The ion explosion spike mechanism for formation of charged particle tracks in solids. *J Appl Phys* 36: 3645-3652.
- Fleischer RL, Price PB, Walker RM (1975) *Nuclear Tracks in Solids*. University of California Press, Berkeley.
- Fletcher JM, Kohn BP, Foster DA, Gleadow AJW (2000) Heterogeneous Neogene cooling and uplift of the Los Cabos block, southern Baja California: Evidence from fission track thermochronology. *Geology* 28: 107-110.
- Foster DA, Gleadow AJW (1992a) Reactivated tectonic boundaries and implications for the reconstruction of southeastern Australia and northern Victoria Land, Antarctica. *Geology* 20: 267-270.
- Foster DA, Gleadow AJW (1992b) The morphotectonic evolution of rift-margin mountains in central Kenya: constraints from apatite fission track analyses. *Earth Planet Sci Lett* 113: 157-171.
- Foster DA, Gleadow AJW (1993) Episodic denudation in East Africa - a legacy of intracontinental tectonism. *Geophys Res Lett* 20: 2395-2398.
- Foster DA, Gleadow AJW (1996) Structural framework and denudation history of the flanks of the Kenya and Anza Rifts, East Africa. *Tectonics* 15: 258-271.
- Foster DA, John BE (1999) Quantifying tectonic exhumation in an extensional orogen with thermochronology: Examples from the southern Basin and Range Province. In: Ring U, Brandon MT, Lister G, Willett SD (eds.) *Exhumation Processes: Normal Faulting, Ductile Flow, and Erosion*, Geol Soc London Special Publ 154: 343-364.
- Foster DA, Raza A (2002) Low-temperature thermochronological record of exhumation of the Bitterroot metamorphic core complex, northern Cordilleran Orogen. *Tectonophysics* 349: 23-36.
- Foster DA, Miller DS, Miller CF (1991) Tertiary extension in the Old Woman Mountains area, California: evidence from apatite fission track analysis. *Tectonics* 10: 875-886.
- Foster DA, Gleadow AJW, Reynolds SJ, Fitzgerald PG (1993) The denudation of metamorphic core complexes and the reconstruction of the Transition Zone, west-central Arizona: constraints from apatite fission-track thermochronology. *J Geophys Res* 98: 2167-2185.
- Foster DA, Gleadow AJW, Mortimer G (1994) Rapid Pliocene exhumation in the Karakoram, revealed by fission-track thermochronology of the K2 gneiss. *Geology* 22: 19-22.
- Fuchs LH (1962) Occurrence of whitlockite in chondritic meteorites. *Science* 137: 425-426.
- Galbraith RF (1990) The radial plot: graphical assessment of spread in ages. *Nucl Tracks Radiat Meas* 17: 207-214.
- Galbraith RF, Green PF (1990) Estimating the component ages in a finite mixture. *Nucl Tracks Radiat Meas* 17: 197-206.
- Galbraith RF, Laslett GM (1993) Statistical models for mixed fission track ages. *Nucl Tracks Radiat Meas* 21: 459-480.
- Gallagher K (1995) Evolving temperature histories from apatite fission-track data. *Earth Planet Sci Lett* 136: 421-435.
- Gallagher K, Brown RW (1997) The onshore record of passive margin evolution. *J Geol Soc London* 154: 451-457.
- Gallagher K, Brown RW (1999a) Denudation and uplift at passive margins: the record on the Atlantic Margin of southern Africa. *Phil Trans Roy Soc London A* 357: 835-859.

39. Naeser CW (1967) The use of apatite and sphene for fission-track age determinations. *Geol Soc America Bull* 78: 1523-1526.
40. O'Sullivan PB, Tagami T (2005) Fundamental of Fission – Track Thermochronology 58: 19-47.
41. Turner DL, Forbes RB, Naeser CW (1973) Radiometric ages of Kodiak Seamount and Giacomini Guyot, Gulf of Alaska: implications for Circum-Pacific tectonics. *Science* 182: 579-581.
42. Turner DL, Frizzell VA, Triplehorn DM, Naeser CW (1983) Radiometric dating for ash partings in coal of the Eocene Puget Group, Washington: Implications for paleobotanical stages. *Geology* 11: 527-531.
43. van der Beek PA (1997) Flank uplift and topography at the central Baikal Rift (SE Siberia): A test of kinematic models for continental extension. *Tectonics* 16: 122-136.
44. van der Beek PA, Cloetingh S, Andriessen PAM (1994) Mechanisms of extensional basin formation and vertical motions at rift flanks: Constraints from tectonic modelling and fission track thermochronology. *Earth Planet Sci Lett* 121: 417-433.
45. van der Beek PA, Andriessen PAM, Cloetingh S (1995) Morpho-tectonic evolution of rifted continental margins: inferences from a coupled tectonic-surface processes model and fission-track thermochronology. *Tectonics* 14: 406-421.
46. van der Beek PA, Delvaux D, Andriessen PAM, Levi KG (1996) Early Cretaceous denudation related to convergent tectonics in the Baikal region, SE Siberia. *J Geol Soc London* 153: 515-523.
47. Villa F, Grivet M, Rebetez M, Dubois C, Chambaudet A, et al. (2000) Damage morphology of Kr tracks in apatite: dependence on thermal annealing. *Nucl Instr Meth B* 168: 72-77.
48. Vineyard GH (1976) Thermal spikes and activated processes. *Radiat Effects* 29: 245-248.
49. Wagner GA (1968) Fission track dating of apatites. *Earth Planet Sci Lett* 4: 411-415.
50. Wagner GA (1969) Traces of the spontaneous fission of ²³⁸Uran as a means of dating of apatite and a contribution to the geochronology of the Odenwald. *N Jahrb Mineral Abh* 110: 252-286.
51. Wagner GA (1977) Fission track dating of apatite and titanite from the Ries: A Contribution to the age and thermal history. *Geol Bavarica* 75: 349-354.
52. Wagner GA, Reimer GM (1972) Fission track tectonics: The tectonic interpretation of fission track apatite ages. *Earth Planet Sci Lett* 14: 263-268.
53. Wagner GA, Storzer D (1970) The interpretation of fission track ages (fission track ages) using the example of natural glasses, apatite and zircon. *Eclogae Geol Helv* 63: 335-344.
54. Wagner GA, Storzer D (1972) Fission track length reductions in minerals and the thermal history of rocks. *Trans Am Nucl Soc* 15: 127-128.
55. Wagner GA, Storzer D (1977) Fission track dating of meteorite impacts. *Meteoritics* 12: 368-369.

Citation: Ghosh AK, Sharma VK, Singh RK (2016) Uplift History of Syenite Rocks of the Sushina Hill, Tamar Porapahar Shear Zone (TPSZ), Purulia: Constraints from Fission-track Ages of Two Cogenetic Minerals. *J Geol Geophys* 5: 245. doi:[10.4172/2381-8719.1000245](https://doi.org/10.4172/2381-8719.1000245)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsgroup.org/journals/submission>